

GRB Redshift Distribution is Consistent with GRB Origin in Evolved Galactic Nuclei

V.I. Dokuchaev¹, Yu.N. Eroshenko¹, and L.M. Ozernoy^{2,3}

¹*Institute for Nuclear Research, Russian Academy of Sciences, Moscow, 117312, Russia*

²*George Mason University, Fairfax, VA 22030-4444, USA*

³*Goddard Space Flight Center, Code 685, Greenbelt, MD 20771, USA*

Abstract. Recently we have elaborated a new cosmological model of gamma-ray burst (GRB) origin [1], which employs the dynamical evolution of central dense stellar clusters in the galactic nuclei. Those clusters inevitably contain a large fraction of compact stellar remnants (CSRs), such as neutron stars (NSs) and stellar mass black holes (BHs), and close encounters between them result in radiative captures into short-living binaries, with subsequent merging of the components, thereby producing GRBs (typically at large distances from the nucleus).

In the present paper, we calculate the redshift distribution of the rate of GRBs produced by close encounters of NSs in distant galactic nuclei. To this end, the following steps are undertaken: (i) we establish a connection between the parameters of the fast evolving central stellar clusters (*i.e.* those for which the time of dynamical evolution exceeds the age of the Universe) with masses of the forming central supermassive black holes (SMBHs) using a dynamical evolution model; (ii) we connect these masses with the inferred mass distributions of SMBHs in the galactic nuclei and the redshift distribution of quasars by assuming a certain ‘Eddington luminosity phase’ in their activity; (iii) we incorporate available observational data on the redshift distribution of quasars as well as a recently found correlation between the masses of galaxies and their central SMBHs. The resulting redshift distribution of the GRB rate, which accounts for both fast and slowly evolving galactic nuclei is consistent with that inferred from the BATSE data if the fraction of fast evolving galactic nuclei is in the range $\varepsilon \simeq 0.016 - 0.16$.

INTRODUCTION

In our previous work [1], we have considered the generation of GRBs due to the radiative collisions of CSRs (for simplicity, taken to be solely NSs) in the central stellar clusters of evolved galactic nuclei. Depending on its initial radius and mass, the characteristic dynamical evolution time of a cluster, t_e , can either exceed the age of the Universe, t_0 , (we call such clusters as ‘*slowly evolving clusters*’ hereinafter) or be less than t_0 (‘*fast evolving clusters*’). The GRB rate from slowly

evolving clusters does not depend on redshift of the host galactic nuclei. Following an approach developed in [2], the mean rate of radiative collisions of NSs in the fiducial galactic nucleus with mass of a central cluster M , radius R , and stellar velocity dispersion $v = (GM/2R)^{1/2}$ is given by [1]:

$$\dot{N}_c \simeq 9\sqrt{2} \left(\frac{v}{c}\right)^{17/7} \frac{c}{R} \simeq 5.8 \cdot 10^{-6} \left(\frac{M}{10^7 M_\odot}\right)^{17/14} \left(\frac{R}{0.1 \text{ pc}}\right)^{-31/14} \frac{\text{events}}{\text{yr galaxy}}, \quad (1)$$

where we use for normalization the likely values of M and R to fit the inferred rate of GRBs. The GRB rate from a unit comoving volume is $\dot{n}_0 = \langle \dot{N}_c \rangle n_g$, where the number density of galaxies $n_g \sim 10^{-2} \text{ Mpc}^{-3}$ and $\langle \dots \rangle$ is an averaging throughout all types of slowly evolving galactic nuclei.

A galactic nucleus with the gravitationally dominating SMBH of mass $M_h > M$ might belong to slowly evolving galactic nuclei as well. The corresponding GRB rate from such a nucleus is given by [1]

$$\dot{N}_{c,h} \simeq \frac{9}{2\sqrt{2}} \left(\frac{Nm}{M_h}\right)^2 \left(\frac{v}{c}\right)^{17/7} \frac{c}{R}, \quad (2)$$

which differs from that given by Eq. (1) by a factor $(M/M_h)^2 \ll 1$.

One more possible contribution to GRB rate from a slowly evolving system is due to the coalescence of tight NS binaries in the galactic discs (which is a commonly believed scheme of cosmological GRB origin). In this paper, we assume that there was no strong evolution with cosmological time of the number of tight NS binaries both in the galactic disks and in the slowly evolving nuclei so that all these contributions result in the redshift-independent GRB rate $\dot{n}_0 \simeq 10^{-8} \text{ yrs}^{-1} \text{ Mpc}^{-3}$.

Meanwhile the actual parameters of galactic nuclei vary in a wide range and some central clusters may, in fact, be fast evolving ($t_e < t_0$). The GRB rate from the fast evolving clusters depends on redshift. Below, we find the redshift distribution of the GRB rate from both slow and fast evolving clusters by (i) modelling the dynamical evolution of a fast evolving cluster in the galactic nucleus, which leads to the production of a SMBH; (ii) connecting the parameters of a fast evolving cluster with the mass of its central SMBH, and (iii) using the observed distributions in luminosity and redshift for quasars, as well as in mass for SMBHs in the galactic nuclei.

FAST EVOLVING GALACTIC NUCLEI

We introduce a dimensionless time variable $y \equiv t_0/t_e = (1+z_e)^{3/2}$, where t_e is the time necessary for a SMBH formation, z_e is the redshift at that instant. Evidently, $y > 1$ for clusters collapsed to the present epoch $t = t_0$ (fast evolving clusters) and $y < 1$ for clusters which do not collapse by t_0 (slowly evolving clusters). As we show elsewhere [3], the dynamical evolution of a stellar cluster, in the framework of a homologous Fokker-Plank approximation [2], leads to the following relationship

between the initial mass of the cluster and the mass of the central SMBH formed there:

$$M_f = 2 \cdot 10^6 y^{2.22} \left[\frac{M_i}{2.8 \cdot 10^8 M_\odot} \right]^{5.44} M_\odot. \quad (3)$$

There are serious observational indications [4], [5], [6] that a fraction of galaxies with a central SMBH might be as high as $\varepsilon \sim 0.1$ and that there is a correlation between the central SMBH mass, M_h , and the luminosity of the host galaxy bulge, L_s , which in a simplified form can be expressed as $M_h \simeq 10^{-2.5}(L_s/L_\odot)M_\odot$. This correlation, taken together with the Schechter luminosity function of galaxies [7], gives the mass distribution of SMBHs, $\phi_1(M_h)dM_h$ with $\phi_1(M_h) \propto \varepsilon$. Then we use the observed [8] distribution of quasars in absolute magnitude M_B and redshift z , $\phi_2(M_B, z)dM_B dz$, for $z \leq 3$. Finally, we assume the existence of an ‘Eddington luminosity phase’ in the evolution of quasars, when they shine with the Eddington luminosity (or with its fraction $\lambda \leq 1$) during a certain finite time interval, which duration only depends upon λ and the initial SMBH mass. In our calculations, the ‘final’ (i.e., by the end of the dynamical evolution) mass of a galactic nucleus is the initial mass of its new-born SMBH. The latter then experiences the ‘Eddington luminosity phase’, during which its mass grows up to the SMBH mass in the actually observed quasar. As a result, we find the mass distribution of *active* SMBHs, $P(M_f, y)dM_f dy$, in terms of the variables y and mass M_f at the instant of the SMBH formation (in the limit $M_f < 3 \cdot 10^6 M_\odot$). Thus, these active SMBHs arise due to the dynamical evolution of galactic nuclei with fast evolving central clusters.

DISTRIBUTION OF GRB RATE IN REDSHIFT

In fast evolving galactic nuclei, the resulting rate of GRB generation per unit of comoving volume is given as a function of redshift z by

$$\dot{n}_{ev}(z) = \int dM_f \int_1^{x(z)^3} P(M_f, y) \dot{N}_{ci}(M_i(M_f), y) \frac{dy x^{3\xi}}{(x^3 - y)^\xi}, \quad (4)$$

where $x = (1 + z)^{1/2}$, $\xi \simeq 0.6$, $\dot{N}_{ci}$ is found from Eq. (1) with $M = M_i$ and $R = R_i$, and function $M_i(M_f)$ is the inverse function given by Eq. (3). Numerical integration over the variable M_f in Eq. (4) proceeds in the range $M_f \sim 10^5 - 3 \cdot 10^6 M_\odot$, which approximately correspond to the range of $M_i \sim 10^7 - 2 \cdot 10^8 M_\odot$.

The total rate of GRBs from *all* kinds of galactic nuclei, $\dot{n}(z)$, includes also the contribution $\dot{n}_0 \simeq 10^{-8} \text{ Mpc}^{-3} \text{ yr}^{-1}$ from all slowly evolving galactic nuclei and disks with $t_e > t_0$, i.e. $\dot{n}(z) = \dot{n}_0 + \dot{n}_{ev}(z)$. The total GRB rate as a function of z can be fitted by a one-parametric function: $\dot{n}(z) = \dot{n}_0(1 + z)^\beta$, where the range of β , $1.5 < \beta < 2$, accounts for the statistical uncertainties [9]. This function is shown in Fig. 1. The consistency between the calculated and inferred GRB rates is achieved at $\varepsilon/\lambda \simeq 0.16$. Since quasars shine at $L \simeq (0.1 - 1) L_{\text{Ed}}$ (e.g., [10]), it implies that $\varepsilon \simeq 0.016 - 0.16$, which is consistent with available data [4]-[6].




FIGURE 1. The GRB rate as a function of redshift. The hatched region confines the allowed GRB redshift distribution from the BATSE data [9]. Three solid lines show the GRB rate from fast evolving stellar clusters in distant galactic nuclei (a constant contribution from the slowly evolving systems is added) as given by our model for the three different values of ϵ/λ ratio, where ϵ is the fraction of galactic nuclei in which SMBHs are formed, and λ is the average quasar luminosity normalized to the Eddington luminosity. For comparison, the dashed line shows the rate of GRBs from NS coalescences in tight binaries produced in the distant star-forming galaxies [11].

REFERENCES

1. Dokuchaev, V.I., Eroshenko, Yu.N., & Ozernoy, L.M., 1998, ApJ, 502, 192
2. Quinlan, G.D. & Shapiro, S.L., 1987, ApJ, 321, 199
3. Dokuchaev, V.I., Eroshenko, Yu. N., & Ozernoy, L.M., 1998, in preparation
4. Van Der Marel, R.P., 1997, astro-ph/9712076
5. Magorrian, J. et al., 1997, submitted to AJ, astro-ph/9708072
6. Kormendy, J. & Richstone, D., 1995, ARA&A, 33, 581
7. Efstathiou, G., Ellis, R.S., & Peterson, B.A., 1988, MNRAS, 232, 431
8. Boyle, B.J., 1991, in “Proc. Texas/ESO–CERN Symposium on Relativistic Astrophysics, Cosmology, Fundamental Physics”, ed. J. D. Barrow et al., p.14
9. Horack J.M. et al., 1995, ApJ, 447, 474
10. Wandel, A., 1998, astro-ph/9808171
11. Bagot, P. et al., 1998, astro-ph/9802094